



Washington State Department of Agriculture • Registration & Services Branch  
1111 Washington Street • PO Box 42589 • Olympia, WA 98504-2589 • Phone: (360) 902-2030 • Fax: (360) 902-2093

## Section 18 Label Approval

No. of Pages including this cover letter: 2 Date: June 6, 2002

<b>Product:</b> tebuconazole (Folicur 3.6F Foliar Fungicide) <b>Site:</b> hops	<b>File Symbol:</b> 02-WA-19
---	------------------------------

Attached is a copy of the approved Section 18 label for the use of **tebuconazole (Folicur 3.6F Foliar Fungicide, EPA Reg No. 3125-394) for control of powdery mildew on hops**. Provisions include the rate, method of application, total number of acres to be treated, use period and other applicable information.

Please Note: These documents are for information purposes only. A copy of the approved label must be obtained from Bayer Corporation prior to using **Folicur 3.6F Foliar Fungicide under this specific exemption**.

### IMPORTANT:

- **THIS EXEMPTION BECOMES EFFECTIVE ON JUNE 15, 2002 AND WILL EXPIRE ON SEPTEMBER 22, 2002.**
- **DO NOT USE THIS PRODUCT ON HOPS UNDER THIS SECTION 18 BEFORE JUNE 15, 2002 OR AFTER SEPTEMBER 22, 2002.**
- **DO NOT USE MORE THAN TWO APPLICATIONS OR MORE THAN A TOTAL OF 16 FLUID OUNCES OF FOLICUR 3.6F FOLIAR FUNGICIDE PER ACRE PER YEAR.**

**Applicators must follow all requirements of the Section 18 label.**

If you would like a copy mailed to you, please contact this office at (360) 902-2030.

SLF:r

# Section 18 Emergency Exemption

SUPPLEMENTAL LABELING FOR DISTRIBUTION AND USE ONLY IN WASHINGTON

## Folicur<sup>®</sup> 3.6 F

### Foliar Fungicide

EPA Reg. No. 3125-394

EPA File Symbol: 02-WA-19

For control of powdery mildew infesting hops.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

All applicable directions, restrictions, and precautions on EPA-registered product label for FOLICUR 3.6F must be followed, unless otherwise superseded by this document.

#### ENVIRONMENTAL HAZARDS

This pesticide is toxic to freshwater, estuarine and marine fish and invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

CROP	DISEASE	RATE OF FOLICUR 3.6F
Hops	Powdery mildew ( <i>Sphaerotheca macularis</i> )	6 to 8 fl oz per acre per application

Apply FOLICUR 3.6F as a preventative spray. The recommended timing is before bloom or after cones are nearing mature size. FOLICUR 3.6F should not be used during bloom due to the potential for impact on cone size and weight.

There must be a minimum 14-day interval between applications. Apply the high rate to highly susceptible hop varieties and/or under severe disease conditions. Apply by aerial or ground equipment in sufficient water for thorough coverage. Increase spray volume as vine growth increases.

**GENERAL DIRECTIONS:** FOLICUR may be applied up to 14 days before harvest. Use of the lowest label rate of an approved surfactant may enhance disease control.

The Washington State Department of Agriculture must be immediately informed of any adverse effects resulting from the use of this pesticide.

**RESTRICTIONS:**

Do not apply more than two (2) applications per acre.

Do not apply more than 16 fl oz of FOLICUR 3.6F per acre per crop season.

**RESISTANCE MANAGEMENT STRATEGY:** FOLICUR is a sterol demethylation inhibitor (DMI) fungicide. Do not mix FOLICUR with another DMI fungicide.

Do not apply through any type of irrigation system.

Do not apply FOLICUR 3.6 F by ground or air within 100 feet of aquatic areas, such as lakes, reservoirs, rivers, permanent streams, marshes or natural ponds, and estuaries.

Do not graze cover crops treated with tebuconazole. Do not harvest cover crops for silage or hay.

This labeling must be in the possession of the user at time of pesticide application.

Effective Date: June 15, 2002

Expiration Date: September 22, 2002

WAFOEM02\_027 06/03/02

Supersedes WAFOEM01\_009

Bayer Corporation  
Crop Protection  
Box 4913  
Kansas City, MO 64120-0013  
(800) 842-8020  
<http://uscrop.bayer.com>

#### IMPORTANT

Before using this product, read and carefully observe the directions, cautionary statements and other information appearing on the product packaging label. This product is sold subject to the Conditions of Sale set forth on the container label.

