

Section 18 Emergency Exemption for the use of Gustafson LSP on Lentil Seed
in the State of Washington only

For use and distribution only within the following counties of Washington, Adams, Asotin, Benton, Columbia, Franklin, Garfield, Grant, Kittitas, Klickitat, Lincoln, Okanogan, Pend Orielle, Stevens, Spokane, Walla Walla and Whitman.
Effective: April 16, 2002 to June 1, 2002

Gustafson LSP Flowable Fungicide
EPA Reg. No. 134
EPA File Symbol 02-WA-14

Active Ingredient:	
Thiabendazole (CAS No. 148-79-8).....	30.0%
Other Ingredients:.....	70.0%
Total:.....	100.0%

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Follow all applicable directions, restrictions, Worker Protection Standard requirements and precautions on the EPA-registered Gustafson LSP Fungicide label.

This label must be in the possession of the user at the time of pesticide application.

Seed treated with Gustafson LSP Fungicide will not provide postemergent protection of plants grown in or near infected fields.

Lentils - To aid in control of seedborne Ascochyta Blight: For ready-mix or slurry treaters, mix 1.7 - 3.0 fl. oz. of Gustafson LSP Flowable Fungicide in an equal amount of water (1:1) to provide adequate coverage and apply to 100 pounds of seed (375 - 750 ppm). Keep dispersion under constant agitation.

The following use restrictions must be followed: a) When handling thiabendazole, or thiabendazole treated seed, work in a well ventilated area and wear chemical resistant gloves; b) Protective eyewear must be worn when handling thiabendazole; c) Do not mix with bare hands; d) Keep out of reach of children; e) Use good personal hygiene, washing hands and exposed skin before eating, drinking, chewing gum, using tobacco, or using the toilet; no food, drink, or tobacco may be allowed in areas of chemical storage or use; f) May be harmful if swallowed or absorbed through skin; wash exposed skin thoroughly with soap and water after use. g) Remove clothing immediately if pesticide gets inside; then wash thoroughly and put on clean clothing.

Manual seed treatment is prohibited under this exemption.

This product is toxic to fish. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Treated seed may be toxic to birds and other wildlife. Clean up any spilled seed immediately, and ensure that treated seed is properly incorporated at planting.

Treated seed must not be used for food, feed, or oil processing (left-over treated seed should not be stored but should be double-sown around the headland).

Do not graze or feed livestock on treated areas for four weeks after planting.

Seed Labeling: Treated seed must be labeled in accordance with the U.S Seed Act as follows:

This seed treated with Thiabendazole at the manufacturer's recommended rate. Do not use for food, feed or oil purposes.

Use with an EPA - approved dye which imparts an un - natural color to the seed.

Gustafson LSP manufactured by:
Gustafson LLC
1400 Preston Road, Suite 400
Plano, Texas 75093
Questions? (972) 985-8877