



Section 18 Emergency Exemption

FOR DISTRIBUTION AND USE ON CHICKPEA IN THE STATE OF WASHINGTON ONLY
DURING 2002 EMERGENCY EXEMPTION PURSUANT TO SECTION 18 OF FIFRA, AS
AMENDED

QUADRIS⁰ Flowable Fungicide

EPA File Symbol No. 02-WA-18

Effective July 1 – September 30, 2002

For Control Of Ascochyta Blight In Chickpeas

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

FAILURE TO FOLLOW THE USE DIRECTIONS AND PRECAUTIONS ON THIS LABEL MAY RESULT IN PLANT INJURY OR POOR DISEASE CONTROL.

INTEGRATED PEST (DISEASE) MANAGEMENT: QUADRIS Flowable Fungicide should be integrated into an overall disease management strategy for each field. Cultural and agronomic practices known to reduce disease development should be followed. These include adapted least susceptible variety, optimum plant population, proper fertilization, crop rotations and water management practices. Consult your local agricultural authorities for additional IPM strategies established for your area.

RESISTANCE MANAGEMENT: A disease management program that includes alternation between QUADRIS Flowable Fungicide and other labeled fungicides that have a different mode of action is essential to prevent pathogen populations from developing resistance to QUADRIS Flowable Fungicide.

CROP	TARGET PEST	RATE		APPLICATION DIRECTIONS
		Fl. oz/A	lb ai/A	
Chickpea	Ascochyta blight (Ascochyta rabei)	6.2 – 9.2 fl oz/A	0.1 – 0.15 lb ai/A	Apply when conditions favor disease. Make the first application at the onset of symptoms or at early flowering if conditions warrant. A second application may be made a minimum of 7 – 14 days later. Use higher rates under severe disease pressure. Applications may be made by ground or air. An adjuvant may be added at recommended rates to improve coverage. Attention: QUADRIS Flowable Fungicide is extremely phytotoxic to certain apple varieties.

RESTRICTIONS:

Do Not apply Quadris within one day of harvest.

Maximum of 2 applications per season.

Do Not graze or harvest treated chickpeas for forage.

Do Not harvest hay from treated fields for a minimum of 21 days following application.

Do Not apply more than 18.4 fl. oz. of Quadris Flowable Fungicide (0.40 lbs a.i.) per acre per crop.

ATTENTION

AVOID SPRAY DRIFT. Extreme care must be used to prevent injury to apple trees (and apple fruit).

DO NOT spray **QUADRIS Flowable Fungicide** where spray drift may reach apple trees.

DO NOT spray when conditions favor drift beyond area intended for application. Conditions which may contribute to drift include thermal inversion, wind speed and direction, sprayer nozzle/pressure combinations, spray droplet size, etc. Contact your State extension agent for spray drift prevention guidelines in your area.

DO NOT use spray equipment that has been previously used to apply **QUADRIS Flowable Fungicide** to spray apple trees. Even trace amounts can cause unacceptable phytotoxicity.

AVOIDING SPRAY DRIFT IS THE RESPONSIBILITY OF THE APPLICATOR.

All applicable directions, restrictions, precautions and environmental hazard statements on the Quadris (EPA Reg. No. 10182-415) container label are to be followed.

This labeling must be in the possession of the user at the time of application.

Any adverse effects resulting from the use of Quadris under this emergency exemption must be immediately reported to the Washington State Department of Agriculture.

QUADRIS® is a trademark of Syngenta Crop Protection, Inc. (legacy Company formed by the merger of Zeneca AG products, Inc. and Novartis Crop Protection, Inc.)

Registrant:
Zeneca AG Products, Inc.
Greensboro, NC 27419

WA1098039CB0702

FN: C:WAQuadrisChickpeas02Sec18Lbl 7-16-02