

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Rally* 40W Fungicide

Rohm and Haas EPA Reg. No. 707-221
Dow AgroSciences EPA Reg. No. 62719-411

This supplemental labeling may be used with either
Dow AgroSciences or Rohm and Haas labeled products.

Section 18 Specific Exemption

For Distribution and Use Only Within Benton and Yakima
Counties in the State of Washington

For Control of Powdery Mildew in Hops

- **This Section 18 Specific Exemption is Valid from March 25, 2002 to September 22, 2002.**
- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for Rally* 40W fungicide before applying. Carefully follow all precautionary statements and applicable use directions.
- In addition to restrictions and limitations imposed by this supplemental labeling, use of Rally 40W is subject to all use precautions and limitations imposed by the label affixed to the product container.

Directions for Use:

Apply 2 to 10 oz. of Rally 40W fungicide per application per acre using ground or aerial equipment. Apply by ground in sufficient water for thorough coverage or by air in a minimum of 10 gallons of water per acre.

Emergence to training: Allow an interval of 7-10 days between applications of Rally 40W fungicide. Use the shorter spray interval on susceptible varieties or under heavy disease pressure. Growers should apply 2 to 4 ounces of Rally 40W per application per acre at this growth stage, but not less than 2 ounces of this product per acre or adequate efficacy may not be achieved.

Training to wire: Allow a spray interval of 5-10 days between treatments of Rally 40W fungicide prior to the beginning of bloom when hop vines are rapidly growing. Use the shorter spray interval on susceptible varieties or under heavy disease pressure. Growers should apply 4 to 6 ounces Rally 40W fungicide per application per acre at this growth stage, but not less than 4 ounces of this product per acre or adequate efficacy may not be achieved.

Wire to 14 days pre-harvest: Apply Rally 40W fungicide at an interval of 7-10 days between applications of Rally 40W fungicide. Use the shorter spray interval on susceptible varieties or under heavy disease pressure. Growers should apply 6 to 10 ounces Rally 40W fungicide per application per acre at this growth stage, but not less than 6 ounces of this product per acre or adequate efficacy may not be achieved.

Restrictions :

- **Do not** apply more than 8 applications per acre per season.
- **Do not** apply more than 5 pounds of Rally 40W (2 lbs active) per acre per season.
- **Do not** enter or allow worker entry into treated areas during the restricted-entry interval (REI) of **48** hours.
- **Do not** apply within 14 days of harvest.
- **Do not** graze livestock in treated areas or harvest crops grown in treated areas for silage or hay.
- **Chemigation: Do not** apply Rally 40W fungicide through any type of irrigation system.

Environmental Hazards:

Do not apply directly to water, or to areas below the mean high-water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Do not apply when weather conditions favor drift or runoff from areas treated. The Washington State Department of Agriculture must be immediately informed of any adverse effects resulting from the use of this pesticide.

*Trademark of Dow AgroSciences LLC

D06-867-001
EPA File Symbol: 02-WA-07
Accepted: 03/22/2002
Initial DAS printing.

Revisions: Renewal of Section 18 exemption for 2002.