



Section 18 Emergency Exemption

SECTION 18 EMERGENCY EXEMPTION FOR THE USE OF SWITCH™62.5WG ON CANEBERRIES IN THE STATE OF WASHINGTON

EFFECTIVE: MAY 1, 2002

EXPIRES: JULY 15, 2002

SWITCH 62.5WG™

EPA FILE SYMBOL NOS. 02-WA-11 (FLUDIOXONIL), 02-WA-10 (CYPRODINIL)

For control of gray mold (*Botrytis cinerea*) in Caneberries

Active Ingredient:	
Cyprodinil: (CAS No. 121552-61-2)	37.5%
Fludioxonil: (CAS No. 131341-86-1)	25.0%
<hr/>	
Other Ingredients:	37.5%
<hr/>	
Total:	100.0%

KEEP OUT OF REACH OF CHILDREN

CAUTION

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND ALL PRECAUTIONS ON THIS LABEL MAY RESULT IN CROP INJURY, POOR DISEASE CONTROL, OR ILLEGAL RESIDUES.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. This label should be in the possession of the user at the time of application.

Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA-approved Switch 62.5WG container label (EPA Reg No. 100-953).

General Information

Caneberries

In caneberries, Switch 62.5WG controls gray mold (*Botrytis cinerea*). Apply 14 oz. of Switch 62.5WG (0.219 pounds fludioxonil + 0.328 pounds cyprodinil) per application per acre in adequate water (100 – 400 gallons is recommended) using ground application equipment only. The first application is recommended at early bloom (by 5% bloom) and repeat 7 to 10 days later depending upon weather.

Restrictions and Conditions: (1) DO NOT apply more than two consecutive applications of Switch 62.5 WG to deter the onset of resistant strains of the fungus. (2) DO NOT apply more than three applications of Switch 62.5WG per acre per year. (3) DO NOT apply more than 42 ounces of Switch 62.5WG per acre per year. (4) DO NOT enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. (5) DO NOT apply Switch by air or through any type of irrigation system. (6) Any adverse effects, which may result from the use of Switch 62.5 WG on caneberries must be reported to the Washington State Department of Agriculture (7) DO NOT apply directly to water, or where surface water is present, or to intertidal areas below the mean high-water mark. Drift and run-off may be hazardous to aquatic organisms in neighboring areas.

Observe the following precautions when spraying in the vicinity of aquatic areas such as lakes; reservoirs; rivers; permanent streams; marshes or natural ponds; estuaries and commercial fish ponds:

- DO NOT apply within 75-feet (25-yard) of bodies of water such as lakes, reservoirs, rivers, permanent streams, natural ponds, marshes or estuaries.
- For all plantings within 150 feet of bodies of water as described above, spray crops from outside the planting away from the bodies of water.
- Shut off the sprayer when at row ends.
- Spray last three rows windward of aquatic areas using nozzles on one side only, with spray directed away from aquatic areas.
- DO NOT cultivate within 10 feet of aquatic areas as to allow a vegetative filter strip.
- DO NOT apply when weather conditions favor drift to aquatic areas.
- DO NOT apply when gusts or sustained winds exceed 10 mph.
- DO NOT apply during a temperature inversion. Mist or fog may indicate the presence of an inversion in humid areas.

Switch™62.5WG trademark of Syngenta
© 2002 Syngenta

Syngenta Crop Protection, Inc.
Greensboro, NC 27419

WA0953034BB0602
FN:WASWITCHCANEERRIES02Sec18lbl 6/26/02