

FRONT PANEL

**SECTION 18 EMERGENCY EXEMPTION FOR THE USE OF WESTERN POPLAR CLEARWING MOTH DISPENSER ON HYBRID POPLAR IN WASHINGTON**

**FOR DISTRIBUTION AND USE ON HYBRID POPLAR IN WASHINGTON, FOR CONTROL OF WESTERN POPLAR CLEARWING MOTH**

## Western Poplar Clearwing Moth Dispenser

**EPA File Symbols 04-WA-04 and 04-WA-05**

**This exemption valid from April 1 to October 1, 2004**

**Active Ingredients**

E,Z-3,13-Octadecadienol 1.38%

Z,Z-3,13-Octadecadienol 0.35%

**Inert Ingredients** 98.27%

TOTAL 100.00%

Net Weight: 40 mg active ingredients  
per dispenser.

**A BIOCHEMICAL PESTICIDE FOR  
MATING DISRUPTION OF THE  
WESTERN POPLAR CLEARWING MOTH (*Paranthrene robiniae*)**

Keep Out Of Reach Of Children

### **CAUTION**

#### **PRECAUTIONARY STATEMENTS**

#### **HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

Caution: Harmful if absorbed through the skin or inhaled. Causes moderate eye and skin irritation. Avoid contact with skin, eyes and clothing. Avoid breathing vapor. Remove contaminated clothing and wash before reuse. Wash thoroughly with soap and water after handling.

#### **FIRST AID**

**IF Inhaled:**

- Move person to fresh air.
- If person is not breathing call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.
- Call poison control center or doctor for treatment advice.

**IF On Skin Or Clothing:**

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call poison control center or doctor for treatment advice.

**IF In Eyes:**

- Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call poison control center or doctor for advice.

**HOT LINE NUMBER**

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-866-326-6737 for emergency medical treatment information.

FRONT PANEL

WPCM

**ENVIRONMENTAL HAZARDS**

Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment washwaters.

Dispenser  
Manufactured by:

SUTERRA LLC  
213 SW Columbia Street  
Bend, Oregon 97702  
1/800-367-8727  
FAX (541) 388-3705

**EPA Establishment No. 56336-OR-1**

**Net Contents: xxx  
Dispensers**

**DIRECTIONS FOR USE**

**It is a violation of Federal Law to use this product  
in a manner inconsistent with its labeling**

**This labeling must be in possession of the user at the time of application**

**Western Poplar Clearwing Moth Dispenser** is a behavior modifying biochemical product (pheromone) which functions by mating disruption. It is to be used for control of the Western Poplar Clearwing Moth (*Paranthrene robiniae*).

**For Application in Newly Planted Trees (less than 5M height)**

**METHOD OF APPLICATION:** For use on hybrid poplars where Western Poplar Clearwing Moths is a pest. The rate of application is 100-400 dispensers per acre (250 to 600 per hectare). **Do not exceed 150 grams a.i./acre/year.** Apply dispensers by hand to thumb-size limbs to secure placement. Place dispensers in the upper one-third of the tree, or higher. If there is a persistent and prevalent wind, apply one dispenser to every tree on the upwind side of the arbor and then evenly distribute the remaining dispensers in the block.

Wash hands after application. Do not handle other types of insect lures before or after lure placement without thoroughly washing hands. Place monitoring traps in the arbor before handling. Monitor treatments carefully.

Reseal containers immediately after use. Store unused product in a refrigerator or freezer not used for food storage. Do not use from damaged, punctured or unsealed containers.

**APPLICATION NOTES:** For best results apply at the time or shortly before the western poplar clearwing moths begin to emerge in the spring (biofix), or when the moths are first detected in pheromone traps such as BioLure®/Scenturion. This product only affects adult male moths and will have no effect on female moths, eggs or larvae. If application occurs following biofix or during the growing season, the arbor must be treated with insecticide treatments that will effectively control hatching larvae until egg laying by previously mated female moths has ceased to occur. Supplemental insecticide applications are to be based on trap monitoring, field scouting and appropriate degree-day models for local growing area and must be timed to control emerging larvae. Follow the **BioLure Technical Bulletin** for trap placement. Monitor insect infestation with traps and by visual inspection of trees and fruit. Re-apply as needed, based on monitoring results and field scouting.

**STORAGE AND DISPOSAL**

**Do not contaminate water, food or feed by storage or disposal**

**PESTICIDE STORAGE:** Store container in cool place until used. **PESTICIDE DISPOSAL:** Wastes resulting from use of this product may be disposed of on-site or at an approved disposal facility.

**CONTAINER DISPOSAL:** Do not reuse container. Discard in trash.

Expiration Date: October 1, 2004

**WARRANTY AND LIMITATION OF DAMAGES**

Suterra LLC warrants that this material conforms to the chemical description on the label. Suterra LLC neither makes, nor authorizes any agent or representative to make, any other warranty of fitness or of merchantability, guarantee or representation, express or implied, concerning this material. To the extent allowable by law, Suterra LLC's liability for breach of this warranty shall not exceed the purchase price of this product. Buyer and user acknowledge and assume all risks and liabilities resulting from the handling, storage and use of this material.