

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Rally* 40W Fungicide

Section 18 Specific Exemption

For Distribution and Use Only Within Benton and Yakima
Counties in the State of Washington

For Control of Powdery Mildew in Hops

- **Section 18 Specific Exemption:** This label is authorized under provisions of Section 18 of the Federal Insecticide Fungicide and Rodenticide Act as amended.
- **Effective Dates: Applications under this specific exemption may be made from May 1, 2004 until September 1, 2004.**
- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- **All applicable directions, restrictions, and precautions on the registered product label for Rally 40W Fungicide (EPA Reg. No. 707-221 or EPA Reg. No. 62719-411) are to be followed.**

Directions for Use

Apply 2 to 10 oz of Rally* 40W fungicide per application per acre using ground or aerial equipment. Thorough coverage is essential. Apply in sufficient water for thorough coverage using ground equipment or by air in a minimum spray volume of 10 gallons of water per acre.

Emergence to training: Allow an interval of 7-10 days between applications of Rally 40W. Use the shorter spray interval on susceptible varieties or under heavy disease pressure. Apply 2 to 4 oz of Rally 40W per application per acre at this growth stage, but not less than 2 oz of this product per acre or adequate efficacy may not be achieved.

Training to wire: Allow a spray interval of 5-10 days between treatments of Rally 40W prior to the beginning of bloom when hop vines are rapidly growing. Use the shorter spray interval on susceptible varieties or under heavy disease pressure. Apply 4 to 6 oz Rally 40W per application per acre at this growth stage, but not less than 4 oz of this product per acre or adequate efficacy may not be achieved.

Wire to 14 days pre-harvest: Apply Rally 40W at an interval of 7-10 days between applications. Use the shorter spray interval on susceptible varieties or under heavy disease pressure. Apply 6 to 10 oz Rally 40W per application per acre at this growth stage, but not less than 6 oz of this product per acre or adequate efficacy may not be achieved.

Restrictions:

- **Do not** make more than 4 applications for Rally 40W per acre per season.
- **Do not** apply more than 2.5 lb of Rally 40W (1 lb active ingredient) per acre per season.
- **Do not** enter or allow worker entry into treated areas during the restricted-entry interval (REI) of **48** hours.
- **Do not** apply within 14 days of harvest.
- **Do not** graze livestock in treated areas or harvest crops grown in treated areas for silage or hay.
- **Chemigation: Do not** apply Rally 40W fungicide through any type of irrigation system.

- **Resistance Management:** Do not make more than 2 consecutive applications of Rally 40W (triazole) without alternating, for at least 1 application, to a fungicide with a different mode of action.

Environmental Hazards:

Do not apply directly to water, or to areas below the mean high-water mark. Do not contaminate water when disposing of equipment washwater or rinsates. Do not apply when weather conditions favor drift or runoff from areas treated.

Any adverse effects resulting from the use of Rally 40W under this emergency exemption must be immediately reported to the Washington State Department of Agriculture.

*Trademark of Dow AgroSciences LLC
R867-001
File Symbol: 04-WA-07
Accepted: 03/12/04
Replaces D06-867-005

Revisions: Renewal of Section 18 use in the state of Washington for 2004.