

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

Indar* 75WSP

Section 18 Specific Exemption

For Control of Mummy Berry Disease in Blueberries

For Distribution and use only in the following counties within the State of Washington: All counties where highbush blueberries are grown west of the Cascade Mountains (Clark, Skagit, Thurston and Whatcom, Clallam, Cowlitz, Grays Harbor Island, Jefferson, King, Kitsap, Lewis, Pacific, Pierce, San Juan, and Snohomish).

ATTENTION

- **Use Period: March 1, 2004 – June 10, 2004.**
- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- The product Indar* 75WSP fungicide (EPA Reg. No. 62719-421), containing 75% fenbuconazole, manufactured by Dow AgroSciences LLC may be used. Existing stocks of Indar 75WSP purchased from the previous registrant, Rohm and Haas (EPA Reg. No. 707-239) may be used.
- All applicable conditions and restrictions on the federally registered product label for Indar 75WSP as well as the Section 18 use directions must be observed.

Directions for Use

Apply 2 ounces of Indar 75WSP fungicide (1.5 ounces active ingredient) per acre using ground or aerial equipment. Make the first application at bud break (when there is 1/16 inch of new leaf growth on the most advanced vegetative bud) and repeat on a 10-14 day interval. A maximum of 3 applications per year is allowed.

A maximum of five (5) fungicide applications are recommended by Washington State University (WSU) to control mummy berry disease [up to three (3) applications between bud break and bloom to protect new growth from infection by ascospores of the fungus (primary stage) or two (2) applications during bloom to protect flowers from infection by conidiospores (secondary stage)].

For the final two applications during bloom, WSU recommends that growers switch to a mixture of Benlate plus Captan. These applications are made to control both the secondary cycle of mummy berry and botrytis blossom blight.

Restrictions:

- **Chemigation: Do not** apply through any type of irrigation system.
- **Preharvest Interval: Do not** apply within 30 days of harvest.
- **Do not** make more than 3 applications or apply more than 6 ounces (0.28 lb active ingredient) per acre per year.
- **Do not** use any spray adjuvants with Indar 75WSP.
- **Do not** graze livestock in treated fields or offer treated material as a livestock feed item.
- Applications are not permitted within 75 feet of streams, rivers, ponds, lakes, or reservoirs.

Any adverse effects or misuse resulting from the use of Indar 75WSP fungicide in connection with this emergency exemption must be immediately reported to the Washington Department of Agriculture.

*Trademark of Dow AgroSciences LLC

R-845-017

EPA-accepted: 02-24-2004

EPA File Symbol: 04-WA-10